

Vitamin D and Bone Health:

A Practical Clinical Guideline for Management in Children and Young People

Vitamin D and bone health: a practical clinical guideline for management in children and young people

There is currently considerable clinical and academic interest in vitamin D in children and young people. This partly relates to recognition of a resurgence of symptomatic vitamin D deficiency with reports of children presenting with rickets or hypocalcaemic symptoms. An additional development has been the recognition that vitamin D may have a physiological extraskeletal role beyond its traditional function as a key regulator of calcium and bone metabolism. There has been a large increase in requests for measurement of vitamin D with evidence that many individuals have suboptimal vitamin D status, often without symptoms. These guidelines have therefore been produced to guide clinicians in the appropriate investigation and management of vitamin D deficiency in children and young people.

Authors: Paul Arundel and Nick Shaw

Publication date: June 2015

Date for review: June 2018

Funding: The development of this document was funded by the National Osteoporosis Society. We are grateful to the authors for giving their time without charge.

Version: 1

Please send any comments on this practical guide to policy.issues@nos.org.uk

Contents

Glossary and abbreviations	4
Key recommendations	5
The role of vitamin D in paediatric bone health	6
How should we assess vitamin D status?	6
Who should be tested for vitamin D deficiency?	6
Who should be treated?	8
How should vitamin D deficiency be treated?	9
Monitoring	9
Vitamin D toxicity	10
Appendix 1: Guideline on assessing dietary calcium intake	11
References	13

Glossary and abbreviations

C ₃ epimer	3-epi-25(OH)D
25(OH)D calcidiol calcifediol	25-hydroxy vitamin D (calcidiol, calcifediol)
HPLC	high-performance liquid chromatography (linked to either fluorescence or MS (tandem MS))
MS	mass spectrometry
PTH	parathyroid hormone
VDBP	vitamin D binding protein
vitamin D	calciferol (either D ₂ or D ₃)
vitamin D ₂	ergocalciferol
vitamin D ₃	cholecalciferol

Conversion factors

10µg (micrograms) vitamin D = 400IU vitamin D

2.5 nmol/L serum 25OHD = 1 ng/mL serum25OHD

Key recommendations

- Measurement of serum 25(OH)D is the best way to estimate vitamin D status.
- Routine testing of serum 25(OH)D levels is not recommended and should be restricted to children and young people with a clear indication for measurement.
- Primary prevention via safe sunlight exposure, advice regarding dietary sources of vitamin D and multivitamin supplements are recommended for high-risk groups.
- Treatment should consist of oral preparations of vitamin D₂ or D₃ given daily for 8 to 12 weeks.
- Many children with vitamin D deficiency have a poor dietary calcium intake and therefore may also require higher dietary calcium intake or calcium supplementation.

The role of vitamin D in paediatric bone health

Nutritional vitamin D deficiency can lead to health problems in children including rickets, impaired growth, muscle weakness and seizures due to hypocalcaemia.

There is no universal consensus on the biochemical definition of vitamin D deficiency. It is current paediatric practice to use a threshold of serum 25 hydroxyvitamin D (25(OH)D) of less than 25 nmol/L to define vitamin D deficiency. This is based on the fact that symptomatic vitamin D deficiency in children (e.g. rickets, hypocalcaemia) will usually only occur with a serum 25(OH)D level below 25 nmol/L. It is also consistent with current UK and European guidelines.

- Serum 25(OH)D below 25 nmol/L is deficient
- Serum 25(OH)D of 25–50 nmol/L may be inadequate in some people.
- Serum 25(OH)D greater than 50 nmol/L is sufficient for almost the whole population.

How should we assess vitamin D status?

Measurement of serum 25(OH)D is the best way to assess vitamin D status. The assay used should have the ability to recognise all forms of 25(OH)D (D₂ or D₃) equally. In practice this means that it should use either HPLC or, more likely, tandem MS. Tandem MS assays can be subject to interference from metabolites such as the C3 epimer, which is synthesised by babies and young children but has also been detected in adult populations. The role of vitamin D binding protein and the impact on free 25(OH)D concentrations currently remains unclear.

Who should be tested for vitamin D deficiency?

Low levels of vitamin D are common in the UK. It is therefore important to consider whether the child's symptoms or signs could be related to vitamin D deficiency before requesting a vitamin D level.

Routine screening is not recommended.

Indications for testing vitamin D status

- 1) Symptoms and signs of rickets:
 - progressive bowing of legs (bowing of legs can be a normal finding in toddlers)
 - progressive knock knees
 - wrist swelling
 - rachitic rosary (swelling of the costochondral junctions)
 - craniotabes (skull softening with frontal bossing and delayed fontanelle closure)
 - delayed tooth eruption and enamel hypoplasia.
- 2) Other symptoms or conditions associated with vitamin D deficiency:
 - long-standing (> 3 months), unexplained bone pain
 - muscular weakness (e.g. difficulty climbing stairs, waddling gait, difficulty rising from a chair or delayed walking)
 - tetany due to low serum calcium
 - seizures due to low serum calcium (usually in infancy)
 - infantile cardiomyopathy.
- 3) Abnormal investigations:
 - low serum calcium or phosphate, high alkaline phosphatase (greater than the local age-appropriate reference range)
 - radiographs – showing osteopenia, rickets or pathological fractures revealed by radiographs.

4) Chronic disease that may increase risk of vitamin D deficiency:

- chronic renal disease, chronic liver disease
- malabsorption syndromes (e.g. coeliac disease, Crohn's disease, cystic fibrosis).

5) Bone diseases in children where correcting vitamin D deficiency prior to specific treatment would be indicated:

- osteogenesis imperfecta
- idiopathic juvenile osteoporosis
- osteoporosis secondary to glucocorticoids, inflammatory disorders, immobility and other metabolic bone conditions.

In the absence of the above indicators, measurement of vitamin D is not indicated.

Primary prevention

Primary preventative measures (at minimum) should be undertaken in patients at high risk. These include advice about safe sunlight exposure, dietary intake of vitamin D and multivitamin supplements.

Indications for vitamin D supplements

The Department of Health recommends daily vitamin D supplements to the following:

- all children between 6 months and 5 years (unless they are receiving over 500ml of formula milk per day)
- breast-fed infants from the age of 1 month if the mother has not taken vitamin D supplements during her pregnancy

In these cases vitamin D supplements would be delivered through the Healthy Start programme

Other indications for vitamin D supplements:

- Children and young people previously shown to be vitamin D deficient or with a serum 25(OH)D of 25–50 nmol/L should take a supplement containing vitamin D. This should be continued unless there is a significant lifestyle change to improve vitamin D status.

- Vitamin D supplements should be considered in other high-risk groups (see below) for vitamin D deficiency, especially if lifestyle advice is not adhered to.

Groups at high risk of vitamin D deficiency

Children and young people in the following groups are at high risk of vitamin D deficiency. Primary prevention is therefore particularly important for them.

- children and young people with diets insufficient in calcium (see Appendix 1) or with generally poor diets
- exclusively breast-fed babies from the age of 6 months, especially if the mother is also at risk of vitamin D deficiency or the infant has not started to take a good range of solid foods
- exclusively breast-fed babies from 1 month if the mother has not taken vitamin D supplements in pregnancy, or if she is known to be vitamin D deficient or insufficient
- children and young people with limited sun exposure (e.g. veiled and photosensitive patients)
- disabled children and young people who spend very little time outdoors
- children and young people who have darker skin, for example people of African, African-Caribbean or South Asian origin, because their bodies are not able to make as much vitamin D
- children and young people taking anticonvulsants that induce liver enzymes such as phenytoin, carbamazepine, primidone or phenobarbitone
- children and young people with family members with proven vitamin D deficiency.

Season

There is a seasonal variation in vitamin D status in the UK, with lower circulating concentrations seen in the population in winter and late spring, compared to summer and autumn. It may be helpful to take into consideration the likely decline in vitamin D status when determining what to do with a child with a low 25(OH)D concentration in autumn or winter.

Having a low 25(OH)D concentration in late summer may reflect a lifestyle that places the individual at risk of vitamin D deficiency. It is important to state that the physiological significance of a given 25(OH)D concentration at a given moment is the same whatever the time of year.

Dietary vitamin D

Consumption of vitamin-D-rich foods can contribute to improving vitamin D status. Foods rich in vitamin D include:

- oily fish such as sardines, pilchards and mackerel
- eggs, meat and milk (in small and varying amounts)
- margarine, some breakfast cereal, some yoghurt and infant formula, which are fortified with vitamin D.

Who should be treated?

In those patients where 25(OH)D is tested (see the previous section: the results should be acted upon as follows:

- serum 25(OH)D < 25 nmol/L: treatment recommended.
- serum 25(OH)D 25–50 nmol/L:
 - Give advice on safe sun exposure and diet.
 - Advise oral preparations containing vitamin D 400–600 IU/day for patients aged 1 month to 18 years. This should be continued unless there is a significant lifestyle change to improve vitamin D status.
 - Ensure dietary calcium intake is adequate.
 - Retesting is not normally required if the individual is asymptomatic and compliant with multivitamin supplements.

- Serum 25(OH)D > 50 nmol/L: provide reassurance and give advice on maintaining adequate vitamin D status through safe sunlight exposure and diet.

Indications for referral to secondary care

The following circumstances indicate referral to secondary care is warranted:

- repeated low serum calcium concentration with or without symptoms (irritability, brisk reflexes, tetany, seizures or other neurological abnormalities)
 - symptomatic: requires immediate referral to A&E if outpatient
 - asymptomatic: discuss treatment with paediatrician
- underlying complex medical disorders (e.g. liver disease, intestinal malabsorption)
- in children, deformities or abnormalities probably related to rickets
- poor response to treatment despite good adherence (defined as a level of 25(OH)D < 50 nmol/L after 8 to 12 weeks of adherent therapy)
- persisting low serum phosphate or low/high alkaline phosphatase.

How should vitamin D deficiency be treated?

Either Vitamin D₃ or Vitamin D₂

Although there are data to show that vitamin D₃ is more bioavailable than vitamin D₂, both forms have been shown to cure rickets and we would not recommend any difference in the doses used. The doses below are based on what are currently recommended in the *British National Formulary for Children* (BNFC)³. However, these may need to be changed dependent on the availability of other vitamin D preparations and evidence of alternative dosing regimens.

- 1–6 months: 3,000 IU orally daily for 8–12 weeks
- 6 months to 12 years: 6,000 IU orally daily for 8–12 weeks
- 12–18 years: 10,000 IU orally daily for 8–12 weeks; a single or divided oral dose totalling 300,000 units can be considered if there is concern about compliance.

Calcium supplements

Many children with vitamin D deficiency rickets have a poor dietary calcium intake. As their bones are growing, there is a greater risk of negative calcium balance. Therefore, in children consideration should always be given to the need for calcium supplementation. Many children with vitamin D deficiency will have a depleted calcium status and/or a poor calcium intake and may therefore benefit from advice about dietary calcium intake. In some cases calcium supplementation may be worthwhile over the period of vitamin D treatment (see Appendix 1). The recommendations below are from the BNFC:

- birth to 4 years: 0.25 mmol/kg calcium q.d.s
- 5–12 years: 0.2 mmol/kg calcium q.d.s
- 12–18 years: 5–10 mmol calcium q.d.s

Dosing also needs to take into account dietary intake and the size of the child.

There is no place for the use of 1 α -hydroxylated preparations (e.g. alfacalcidol or calcitriol) in the routine management of vitamin D deficiency. Their use is limited to treating significant hypocalcaemia, disorders of malabsorption and renal disease.

Monitoring

Bone profile and vitamin D tests (and a PTH test if the patient has rickets or hypocalcaemia) should be repeated at the end of the course of treatment.

If the 25(OH)D level is greater than 50nmol/L and the bone profile is normal:

- Give advice on safe sun exposure and diet.
- Advise multivitamin containing vitamin D 400–600 IU/day. This should be continued unless there is a significant lifestyle change to improve vitamin D status.

If 25(OH)D is below 50 nmol/L:

- Consider poor compliance, drug interactions and underlying disease such as renal disease, liver disease and malabsorption.
- If poor compliance is suspected, a high-dose treatment may be considered if the patient is aged 12–18 years (e.g. 300,000 IU as a single or divided dose).

Note: If a child's symptoms/signs have not improved despite a satisfactory 25(OH)D concentration, they are unlikely to be related to vitamin D deficiency.

Vitamin D toxicity

Serum 25(OH)D concentrations of above 375nmol/L have been associated with hypercalcaemia and hyperphosphataemia but there is no agreement on the threshold concentration or amount of vitamin D that results in toxicity. This is, in part, a reflection of the paucity of studies that address the safety of vitamin D supplementation. In adults, prolonged daily intake of vitamin D up to 10,000 IU or serum concentrations of 25(OH)D of up to 240 nmol/L appear to be safe. Although described, acute vitamin D intoxication is rare and usually results from vitamin D doses much higher than 10,000 IU per day. However, the long-term effects of supplementation with high doses of vitamin D are not known. Risks such as nephrolithiasis cannot be excluded. Caution is required in any child or young person with a granulomatous disease (e.g. tuberculosis or sarcoidosis).

Appendix 1: Guideline on assessing dietary calcium intake

Dietary reference values for calcium

Age	Reference nutrient intake for calcium mg/day (mmol/day)
0-12 months	525 (13.1)
1-3 years	350 (8.8)
4-6 years	450 (11.3)
7-10 years	550 (13.8)
11-14 years, male	1000 (25.0)
11-14 years, female	800 (20.0)
15-18 years, male	1000 (25.0)
15-18 years, female	800 (20.0)

Note: 1 mmol calcium = 40 mg calcium.

Reference: Department of Health, *Dietary Reference Values for Food, Energy and Nutrients for United Kingdom* (Report 41), London: TSO, 1991.

Appendix 1: Guideline on assessing dietary calcium intake

Calcium-rich foods

Find calcium-rich foods from this list for a bone-healthy diet.^a Serving sizes are based on average portions.^b The details below provide a general guide – precise calcium content can vary depending on source of food stuffs.

Food	Serving size (average)	Calcium (mg)
Milk		
Milk, semi-skimmed	glass, 200 ml	240
Milk skimmed	glass, 200 ml	244
Milk whole	glass, 200 ml	236
Milkshake	takeaway, 300 ml	387
Soy drink, calcium enriched	glass, 200 ml	178
Yoghurt and cream		
Yoghurt, low-fat, fruit	pot, 150 g	210
Yoghurt, low-fat, plain	pot, 150 g	243
Cream, double, whipped	portion, 45 g	26
Cream single	tablespoon, 15 g	13
Cheeses		
Danish blue	portion, 40 g	195
Edam	portion, 40 g	318
Feta	portion, 40 g	144
Camembert	portion, 40 g	94
Cheddar	medium chunk, 40 g	296
Cheese spread	portion, 30 g	149
Cottage	small pot, 112 g	142
Mozzarella, fresh	portion, 56 g	203
Parmesan, fresh	portion, 30 g	308
Vegetables		
Broccoli, boiled	serving, 85 g	34
Watercress, raw	small bunch, 20 g	34
Curly kale	serving, 95 g	143
Okra, stir fried	8 medium, 40 g	88
Red kidney beans, canned	3 tablespoons, 105 g	75
Chick peas, boiled	3 tablespoons, 90 g	41
Green/French beans	serving, 90 g	50
Baked beans	serving, 135 g	72
Nuts		
Almonds	12 whole, 26 g	62
Brazil nuts	6 whole, 20 g	34
Hazelnuts	20 whole, 20 g	28
Sesame seeds	1 tablespoon, 12 g	80
Walnuts	12 halves, 40 g	38
Tahini paste	1 heaped teaspoon, 19 g	129

Food	Serving size (average)	Calcium (mg)
Desserts		
Cheesecake, fruit	slice, 120 g	94
Custard made with milk	portion, 120 g	166
Rice pudding, canned	portion, 200 g	176
Ice cream, dairy, vanilla	serving, 75 g	75
Fromage frais, fruit	small pot, 60 g	52
Fish		
Sardines in oil, tinned	portion, 100 g	500
Whitebait, fried	portion, 80 g	688
Salmon, tinned	portion, 100 g	91
Fish paste	small jar, 35 g	98
Breads and grains		
Pasta, plain, cooked	portion, 230 g	85
Rice, white, boiled	portion, 180 g	32
White bread	slice, 30 g	53
Wholemeal bread	slice, 30 g	32
Muesli, Swiss style	portion, 50 g	55
Fruits		
Apricots, raw, no stone	4 fruit, 160 g	117
Figs, ready to eat	4 fruit, 220 g	506
Currants	2 tablespoons, 50 g	47
Orange	peeled, 160 g	75
Other foods		
Tofu, soy bean, steamed	100 g	510
Omelette, cheese	2 eggs, 120 g	344
Quiche, cheese and egg	slice, 140 g	367
Macaroni cheese	portion, 220 g	374
Pizza, cheese and tomato	9"-10" pizza, 410 g	873
Lasagne	portion, 420 g	420

References

^a Food Standards Agency. *McCance and Widdowson's The Composition of Foods*, Sixth Summary Edition. Cambridge: Royal Society of Chemistry, 2002

^b Food Standards Agency. *Food Portion Sizes 2002*.

Reproduced with permission from Chris Rudd, Registered Dietitian (Dietetic Advisor, Medicines Management Team, NHS Sheffield Clinical Commissioning Group).

References

1. Braegger C, Campoy C, Colomb V et al. Vitamin D in the healthy European paediatric population. *JPGN* 2013;56:692–701.
2. Scientific Advisory Committee on Nutrition. *Update on Vitamin D*. London: TSO, 2007.
3. *British National Formulary for Children 2013–2014*. London: BMJ Group, 2014. Section 9.6.4 Vitamin D.
4. Shaw NJ, Mughal MZ. Vitamin D and child health part 1 (skeletal aspects). *Arch Dis Child* 2013;98:363–367.
5. Shaw NJ, Mughal MZ. Vitamin D and child health part 2 (Extraskeletal and other aspects). *Arch Dis Child* 2013;98:368–372.
6. Aggarwal V, Seth A, Aneja S, Sharma B, Sonkar P, Singh S, Marwaha RK. Role of calcium deficiency in development of nutritional rickets in Indian children: a case control study. *J Clin Endocrinol Metab* 2012;97(10): 3461–3466.
7. Ladhani S, Srinivasan L, Buchanan C, Allgrove J. Presentation of vitamin D deficiency. *Arch Dis Child* 2004;89(8):781–784.
8. Arundel P, Ahmed SF, Allgrove J, Bishop NJ, Burren CP, Jacobs B, Mughal MZ, Offiah AC, Shaw NJ. British Paediatric and Adolescent Bone Group’s position statement on vitamin D deficiency. *BMJ* 2012; 345:e8182.

 **0845 130 3076** (General Enquiries)

 **0845 450 0230** (Helpline)

 **www.nos.org.uk**

 **Camerton, Bath BA2 0PJ**

